arXiv:math/0002131v1 [math.KT] 16 Feb 2000

Chern character for the Schwartz algebra of p-adic $GL(n)$

Jacek Brodzki Roger Plymen

1 Introduction

Atiyah has demonstrated that rational cohomology of a compact Hausdorff space can be defined in terms of K-theory. This is made possible by the existence of a Chern character

$$ch_j : K^j(X) \to H^{ev/odd}(X; \mathbf{Z}), \quad j = 0, 1.$$

As in [1, p. 142](see also [22, Sec. 6.2]), this map is a rational isomorphism:

$$ch_j \otimes_{\mathbf{Z}} 1 : K^j(X) \otimes_{\mathbf{Z}} \mathbf{Q} \simeq H^{ev/odd}(X; \mathbf{Q}). \tag{1.1}$$

When X is a compact smooth manifold, the value of the Chern character on a K-theory class represented by a bundle E can be computed explicitly in terms of an inhomogeneous differential form whose components are (up to factors) exterior powers of the curvature of E. Thus, for instance, information about K-theory of X may be obtained by pairing the cohomology class represented by the Chern character with closed de Rham currents. There is now a construction, due to Gorokhovsky [12], which provides explicit formulae for the classical Chern character of a compact Hausdorff space X in terms of Alexander-Spanier cocycles. In the case when X is a compact smooth manifold, both ways of computing the Chern character coincide.

Connes created a Chern character from K-homology of an algebra A, defined in terms of abstract elliptic operators, with values in cyclic cohomology [4, Chap. I]. He also provided a key idea for the construction of a dual Chern character from the K-theory of A, defined in terms of analogues of vector bundles, with values in cyclic homology [4, Chap. II, Prop. 14]. The two characters are compatible with the index pairing between K-homology and K-theory of the algebra A.

These groundbreaking ideas were developed by a number of authors [14, 15, 17, 19] and completed by Cuntz and Quillen who provided explicit formulae for Chern characters with values in periodic cyclic homology [10]. More recently, Cuntz constructed a universal bivariant kk-functor and a compatible Chern character whose target is the bivariant periodic cyclic cohomology of Cuntz and Quillen [7].

An important application of these constructions is provided by a proof of the Baum-Connes conjecture for the p-adic group $GL(n)$ [2]. There, Chern characters were used to obtain information about the Baum-Connes assembly map. For a locally compact group G the Baum-Connes map is an index map from the universal example of proper G actions to the K-theory of the reduced C^*-algebra $C_r^*(G)$.

In this paper we construct a Chern character

$$ch : K_*(C_r^*(G)) \to HP_*(S(G)) \tag{1.2}$$

where $S(G)$ is the Schwartz algebra of the p-adic group $GL(n)$. We then prove that this map is an isomorphism after tensoring over $\mathbf{Z}$ with $\mathbf{C}$. The first step in our construction is to adapt the formula of Cuntz and Quillen to the context of Fréchet algebras. It turns out that this translation is very natural in the case when the Fréchet algebra under study is a dense subalgebra of a C^*-algebra which is stable under holomorphic functional calculus. The Schwartz algebra $S(G)$ is defined as the strict inductive limit of nuclear Fréchet algebras $S(G//K)$ which satisfy this property [13, p. 93].

Using explicit formulae provided by the connection-curvature computation of the Chern character in the Chern-Weil theory and Gorokhovsky's results in the topological case we compare the map of Cuntz and Quillen with the classical Chern character in algebraic topology. Ultimately our result that the Chern character (1.2) is an isomorphism after tensoring with $\mathbf{C}$ follows from the fact that the classical Chern character is a rational isomorphism as stated in (1.1).

2 Fréchet algebras and the Chern character

Cuntz and Quillen provide explicit formulae [10] for Chern character maps $ch_{CQ} : K_*(A) \to HP_*(A)$ from the K-theory of a unital algebra A to its periodic cyclic homology. The goal of this section is to adapt their results to the context of Fréchet algebras and to show how these maps compare, in the case of the algebra of smooth functions on a manifold, with the Chern character given by the Chern-Weil theory.

By a Fréchet algebra we will understand a locally multiplicatively convex Fréchet algebra over $\mathbf{C}$. That is, a Fréchet algebra is a complete metrizable topological algebra whose topology is given by a countable family of submultiplicative seminorms.

K-theory for Fréchet algebras was constructed by Phillips [20]. Any definition of $K_*(A)$ requires a suitable choice of stabilization of A. In the case of Banach or C^*-algebras one uses infinite matrices over A with finitely many nonzero entries, that is the direct limit $\bigcup_n M_n(A)$ of the natural direct system of matrix algebras $M_n(A)$ of increasing size. This approach is not suitable for a general Fréchet algebra, see [20, Section 2] for a full discussion. It turns out that the stabilization of A best adapted to the definition of K-theory for Fréchet algebras is given by $S(\mathbf{Z}^2) \otimes A$, where $\mathcal{S}(\mathbf{Z}^2)$ is the algebra of rapidly decreasing functions on $\mathbf{Z}^2$. However, there is a significant simplification in the case of algebras which are dense subalgebras of C^*-algebras, stable under holomorphic functional calculus and which are Fréchet algebras in a finer topology. All Fréchet algebras considered in this paper are of this kind unless stated otherwise. In this case Phillips proves that the K-theory can be defined in the following familiar way [20, Def. 7.1, Cor. 7.9]. For a unital Fréchet algebra A, $K_0(A)$ is defined as the Grothendieck group of the semigroup of isomorphism classes of finitely generated projective modules. Thus a class in $K_0(A)$ can be represented by a projection in $M_n(A)$, for some n. Moreover, projections in the class of a projection e are homotopic to e. Let $GL_n(A)$ denote the group of invertible $n \times n$ matrices, and let $GL_n(A)_0$ be the path component of the identity. We define $K_1(A) = \lim GL_n(A)/GL_n(A)_0$. For nonunital algebras $K_i(A)$ is defined as the kernel of the map from $K_i(\tilde{A})$ to $K_i(\mathbf{C})$, where $\tilde{A}$ is the unitization of A.

Cyclic type homology theories of a complex algebra A, for example Hochshild, cyclic and periodic cyclic homology, are computed using a mixed complex $(\Omega A, b, B)$ associated with the algebra A [9]. The universal differential graded algebra ΩA is generated by the algebra A and the symbols da, $a \in A$, which are linear in A and satisfy the Leibniz identity $d(ab) = d(a)b + adb$ for any $a, b \in A$. In degree n, the space of n-forms $\Omega^n A$ is spanned by the elements $a_0 da_1 \dots da_n$. As a vector space, $\Omega^n A = A \otimes (A/\mathbf{C})^{\otimes n}$. The differential $b : \Omega^n A \to \Omega^{n-1} A$ is defined in positive degrees by $b(\omega da) = (-1)^{|\omega|}[\omega, a]$, where $|\omega|$ is the degree of the form ω. In degree zero we put b equal identically zero. The degree $+1$ differential $B : \Omega^n A \to \Omega^{n+1} A$ is defined by $B = \sum_{i=0}^{n} \kappa^i d$, where κ is the degree zero Karoubi operator: $\kappa(\omega da) = (-1)^{|\omega|} da \cdot \omega$. The two operators b and B anticommute and are of square zero:

$$b^2 = Bb + bB = B^2 = 0.$$

For a locally convex algebra A one must choose a suitable topological tensor product to use in the definition of the differential graded algebra ΩA. In this paper we choose the completed inductive tensor product $\bar{\otimes}$ as in [3], to which we refer the reader for further details. Thus, for all $n \geq 0$, $\Omega^n(A) = A \bar{\otimes} (A/\mathbf{C})^{\bar{\otimes} n}$. The differentials b and B are continuous in this topology.

Let A be a unital locally convex algebra over $\mathbf{C}$. By definition, the *Hochschild homology* of A is

$$HH_*(A) = H_*(\Omega A, b).$$

The *periodic cyclic homology* $HP_*(A)$ of A is the homology of the $\mathbf{Z}/2\mathbf{Z}$-graded complex

$$\dots \xrightarrow{B-b} \prod_{n \geq 0} \Omega^{2n}(A) \xrightarrow{B-b} \prod_{n \geq 0} \Omega^{2n+1}(A) \xrightarrow{B-b} \prod_{n \geq 0} \Omega^{2n}(A) \xrightarrow{B-b} \dots$$

Let A be a unital Fréchet algebra. Let $[e] \in K_0(A)$ be the K-class of an idempotent matrix $e \in M_k(A)$ over A. Then the even Chern character of Cuntz and Quillen assigns to $[e]$ an even class in the periodic cyclic homology of A represented by the even periodic cycle

$$ch_{CQ} : e \mapsto \mathrm{Tr} e + \sum_{n \geq 1} \frac{(2n)!}{n!} \mathrm{Tr} \left(\left((e - \frac{1}{2} \right) de^{2n} \right) \in \Omega^{ev} A. \tag{2.1}$$

Here $\Omega^{ev} A = \prod_{n \geq 0} \Omega^{2n} A$.

In the odd case, let us assume that a class $[g] \in K_1(A)$ is represented by an invertible matrix $g \in GL_k(A)$. The odd Chern character maps the class $[g]$ to the odd periodic cyclic homology class represented by the odd cycle

$$ch_{CQ} : g \mapsto \sum_{n \geq 0} n! \mathrm{Tr}(g^{-1}dg(dg^{-1}dg)^n) \tag{2.2}$$

in $\Omega^{odd}(A) = \prod_{n \geq 0} \Omega^{2n+1}A$.

This Chern character is compatible with algebra homomorphisms in the sense that a homomorphisms $\phi : A \to B$ of algebras gives rise to the commutative diagram

$$\begin{tikzcd} K_*(A) \arrow[r, "ch_{CQ}"] \arrow[d, "\phi"] & HP_*(A) \arrow[d, "\phi"] \\ K_*(B) \arrow[r, "ch_{CQ}"] & HP_*(B) \end{tikzcd}$$

This map is compatible with Morita equivalence in the following sense. There is a similar diagram in which B is replaced by the matrix algebra $M_n(A)$ and where vertical arrows are isomorphisms.

The formula of Cuntz and Quillen works for any Fréchet algebra, commutative or not. Its main strength lies in the fact that it can be adapted to coincide with the classical Chern character known from algebraic topology or the Chern character defined for smooth manifolds using the Chern-Weil theory. To make a comparison of the Chern character of Cuntz and Quillen with the one known from the Chern-Weil theory, let us recall the main points in the construction of the latter map.

Let E be a complex vector bundle over a compact smooth manifold M and let ∇ be a connection on E. The curvature R of ∇ is an $\mathrm{End}E$-valued two-form on M. Then $\mathrm{Tr}\exp(R/2\pi i)$ is an inhomogeneous even exact form on M. The component of degree $2k$ represents a class in $H^{2k}_{dR}(M)$. With this understood, the even Chern-Weil character is the map

$$ch_{CW} : [E] \mapsto [\mathrm{Tr}\exp(R/2\pi i)] \in H^{ev}_{dR}(M) = \bigoplus_{n \geq 0} H^{2n}_{dR}(M), \tag{2.3}$$

where $[E] \in K^0(M)$ is the K-theory class represented by the bundle E.

In the odd case, we assume that a class in $K^1(M)$ is represented by an invertible matrix $g \in GL_n(C^\infty(M))$. The odd Chern character is defined by the formula [11, p.491]

$$ch_{CW} : [g] \mapsto \sum_{n\geq 0}\left[(-1)^n \frac{n!}{(2\pi i)^n(2n+1)!}\mathrm{Tr}((g^{-1}dg)^{2n+1})\right] \tag{2.4}$$

which takes values in $H^{odd}_{dR}(M) = \bigoplus_{n\geq 0} H^{2n+1}_{dR}(M)$.

2.5 Remark. The factors $(2\pi i)^{-n}$ in the two formulae above are used in algebraic topology to ensure that the Chern characters represent integral classes in cohomology of M. This plays a crucial role, for instance, in the index theorems of Atiyah and Singer.

When A is the algebra $A = C^\infty(M)$ of smooth functions on a smooth compact manifold M the Cuntz-Quillen maps (2.1) and (2.2) coincide with (2.3) and (2.4) as we now show.

Let us denote by $(\Omega M, d)$ the complexified de Rham complex of the manifold M. The universal property [8] of the differential graded algebra ΩA implies that there is a natural surjection of mixed complexes

$$\mu : (\Omega A, b, B) \to (\Omega M, 0, d)$$

which sends a noncommutative n-form $f^0 df^1 \cdots df^n$ to the differential form $(1/n!) f^0 df^1 \wedge \ldots \wedge df^n$. This map induces an isomorphism of all cyclic type homology theories associated with the two mixed complexes if the algebra A satisfies the Hochschild-Kostant-Rosenberg theorem

$$HH_*(A) = H_*(\Omega A, b) = \Omega M.$$

A crucial result of Connes' shows that this is true for $A = C^\infty(M)$. In particular we have in this case that $HP_{ev/odd}(A) \simeq H^{ev/odd}_{dR}(M)$.

The map μ thus induces a map $\mu : HP_{ev/odd}(A) \to H^{ev/odd}_{dR}(M)$. The de Rham cohomology groups are $\mathbf{C}$-modules with the natural action $\lambda[\omega] = [\lambda\omega]$ for any $\lambda \in \mathbf{C}$. We use this fact to define the scaling map

$$c_n : H^n(M) \to H^n(M)$$

which for $n = 2k$ multiplies by $1/(2\pi i)^k$ and for $n = 2k+1$ by $1/(2\pi i)^{k+1}$. We let $c : \bigoplus_n H^n(M) \to \bigoplus H^n(M)$ be the map whose component in degree n is c_n. It is clear that c is an isomorphism of $\mathbf{C}$-modules. Let $\mu_c = c \circ \mu$.

The comparison of the two Chern characters is established by assembling various known results which we recall here for future reference.

2.6 PROPOSITION. *Let M be a smooth compact manifold and let $A = C^\infty(M)$ be the algebra of smooth functions on M. The following diagram commutes*

$$\begin{tikzcd}
K_*(A) \arrow[r, "ch_{CQ}"] \arrow[d, "SS"] & HP_*(A) \arrow[d, "\mu_c"] \\
K^*(M) \arrow[r, "ch_{CW}"] & H_{dR}^{ev/odd}(M)
\end{tikzcd}$$

where the left vertical arrow is the isomorphism given by the Serre-Swan theorem.

Proof. In the even case, the Serre-Swan map sends a projection $e \in M_k(C^\infty(M))$ to the bundle $E = e\theta^k$, where θ^k is the trivial complex rank-k bundle on M. The trivial bundle θ^k is equipped with the canonical connection d, which is just the de Rham differential. The subbundle E of θ^k is then equipped with the induced Grassmannnian connection

$$D = e \cdot d \cdot e$$

which acts on sections of E. A typical section of E is of the form ef, where $f \in C^\infty(M, \mathbf{C}^k)$ is a section of the trivial bundle θ^k. The curvature of this connection is (cf. [21, p.223]) $R = e(de)^2$. Moreover, $e(de)^2 = (de)^2 e$ so that $R^k = e(de)^{2k}$. Hence the Chern character of the sub-bundle E is

$$ch(E) = \exp(R/2\pi i) = \sum_{n \geq 0} \frac{1}{(2\pi i)^n n!} e(de)^{2n}$$

where the power on the right is taken using the exterior product of forms.

The commutativity of the diagram in the even case is now established using that [10, p. 435]

$$\mu_c ch_{CQ}([e]) = \sum_{n \geq 0} \frac{1}{(2\pi i)^n n!} \mathrm{Tr}(e(de)^{2n}) = ch_{CW}(SS[e])$$

The odd case follows once we notice that [10, p. 436]

$$\mu_c[g] = \sum_{n \geq 0} \frac{n!}{(2\pi i)^n (2n+1)!} \mathrm{Tr}((g^{-1}dg)^{2n+1}).$$

■

3 The Schwartz algebra of $GL(n)$

Let F be a nonarchimedean local field and let $G = GL(n) = GL(n, F)$. Let K be a compact open subgroup of G. We define $S(G//K)$ to be the algebra of all complex-valued functions on G which are rapidly decreasing and K-bi-invariant, with product given by convolution.

3.1 Lemma. *For every compact open subgroup K of G the algebra $S(G//K)$ is an m-convex Fréchet algebra.*

Proof. Mischenko's Fourier transform [18] provides the following isomorphism of locally convex algebras

$$S(G//K) \simeq \bigoplus_M [C^\infty(\mathrm{End}F(M:K))]^{W(M)} \tag{3.2}$$

where $F(M:K) \to X(M:K)$ is a complex, Hermitian, trivialized $W(M)$-bundle over a compact manifold $X(M:K)$. The group $W(M)$ is a finite reflection group and M is a Levi subgroup. One Levi subgroup is chosen in each conjugacy class of G; in particular, the sum on the right is finite. Each direct summand on the right is a subalgebra of the algebra $C^\infty(X(M:K)) \otimes M_n(\mathbf{C})$ of matrix-valued smooth functions on $X(M:K)$. The algebra $C^\infty(V)$ of smooth functions on a compact smooth manifold V is an m-convex Fréchet algebra. Indeed, for every natural number n we put $p_n(f) = \sup\{|\alpha| \le n,\, x \in V \mid |\partial^\alpha f(x)|\}$ and define

$$q_n(f) = \sum_{i=0}^{n} \frac{1}{i!} p_i(f)$$

The seminorms q_n are submultiplicative.

The matrix algebra $M_n(\mathbf{C})$ is equipped with the Hilbert-Schmidt norm $\| \ \|_2$, which is submultiplicative.

The algebraic tensor product $C^\infty(X(M:K)) \otimes M_n(\mathbf{C})$ is topologised by submultiplicative seminorms $q_n \otimes \| \ \|_2$ and so is an m-convex Fréchet algebra. This remark combined with Mischenko's isomorphism shows that $S(G//K)$ is an m-convex Fréchet algebra for every compact open subgroup K of G. ■

Following Harish-Chandra [13, p. 93], the Schwartz algebra of G is defined as the strict inductive limit of the algebras $S(G//K)$, that is

$$S(G) = \varinjlim S(G//K) = \bigcup_K S(G//K).$$

Choose a left-invariant Haar measure on G. Then $L^1(G)$ acts on $L^2(G)$ by convolution

$$\lambda(f)h = f * h,$$

for $f \in L^1(G)$ and $h \in L^2(G)$. The reduced C^*-algebra $C_r^*(G)$ is the closure of $\lambda(L^1(G))$ in the C^*-algebra of bounded operators on $L^2(G)$.

In this section, we prove the following main result of this paper.

3.3 THEOREM. *There exists a Chern character map*

$$ch : K_*(C_r^*(G)) \to HP_*(S(G)) \tag{3.4}$$

which is an isomorphism after tensoring over $\mathbf{Z}$ *with* $\mathbf{C}$.

Proof. For every compact open subgroup K of G the algebra $S(G//K)$ is stable under holomorphic functional calculus, as follows from Mischenko's theorem (3.2). Moreover, each algebra $S(G//K)$ is an m-convex Fréchet algebra by Lemma 3.1. Denote by $C_r^*(G//K)$ the closure of $S(G//K)$ in the reduced C^* algebra $C_r^*(G)$. Then $S(G//K)$ is dense in $C_r^*(G//K)$. Thus the Cuntz-Quillen formula applies to give a Chern character

$$ch_{CQ}^K : K_*(S(G//K)) \to HP_*(S(G//K))$$

This Chern character then determines a map:

$$ch_{CQ}^K : K_*(C_r^*(G//K)) \simeq K_*(S(G//K)) \to HP_*(G//K)$$

The C^*-algebras $C_r^*(G//K)$ form a direct system whose C^*-inductive limit is the reduced C^*-algebra $C_r^*(G)$. We thus have a direct system of Chern characters ch_{CQ}^K and we define

$$ch = \varinjlim ch_{CQ}^K : K_*(C_r^*(G)) \to HP_*(S(G))$$

We use here continuity of K-theory and periodic cyclic homology with respect to direct limits which holds in this case [3, Theorem 6].

We now prove that this character is an isomorphism after tensoring (over $\mathbf{Z}$) with $\mathbf{C}$.

Relying on Mischenko's theorem stated in Lemma 3.1 we proved in [3] that each algebra $S(G//K)$ is Morita equivalent to a commutative algebra

$$S(G//K) \overset{\text{Morita}}{\sim} \bigoplus_M C^\infty(X(M:K))^{W(M)} \tag{3.5}$$

where each direct summand is the algebra of invariant smooth functions on $X(M:K)$.

Using this fact together with the compatibility of functors K_* and HP_* with algebra homomorphisms and Morita equivalence we see that, for every open subgroup K of G, there is the commutative diagram

$$\begin{tikzcd}
K_*(S(G//K)) \arrow[r, "ch^K_{CQ}"] \arrow[d, "\simeq"'] & HP_*(S(G//K)) \arrow[d, "\simeq"'] \\
\bigoplus_M K_*(C^\infty(X(M:K)))^{W(M)} \arrow[r, "ch_{CQ}"] & \bigoplus_M HP_*(C^\infty(X(M:K)^{W(M)}))
\end{tikzcd} \tag{3.6}$$

Vertical arrows in this diagram are isomorphisms. The bottom arrow is the direct sum of Cuntz-Quillen Chern characters defined for each Levi subgroup M using formulae (2.1) and (2.2). The key step in our argument is to show that each of the components of the map at the bottom is an isomorphism after tensoring with $\mathbf{C}$.

This follows from the following geometric consideration. Let X be a compact W-manifold, where W is a finite reflection group. There is the following commutative diagram

$$\begin{tikzcd}
K_*(C^\infty(X)) \arrow[r, "ch_{CQ}"] \arrow[d, equal] & HP_*(C^\infty(X)) \arrow[d, "\mu_c"] \\
K_*(C^\infty(X)) \arrow[r, "ch_{CW}"] \arrow[d, equal] & H^*_{dR}(X) \\
K_*(C^\infty(X)) \arrow[r, "ch_G"] & H^*_{AS}(X). \arrow[u, "\lambda"]
\end{tikzcd} \tag{3.7}$$

The top two rows form the commutative diagram of Proposition 2.6. The bottom row introduces the Chern character with values in the Alexander-Spanier cohomology $H^*_{AS}(X)$, which was constructed by Gorokhovsky for any compact Hausdorff space [12]. Ghorokhovsky proves that his map coincides with the classical Chern character. In particular, it is a rational isomorphism by [16]. Moreover, when X is a compact smooth manifold, there is a natural isomorphism

$$\lambda : H^*_{AS}(X) \to H^*_{dR}(X)$$

which identifies the Alexander-Spanier cohomology with the de Rham cohomogy. It is shown in [12] that this map sends Gorokhovsky's Chern character to the Chern character known from the Chern-Weil theory. In other words, the bottom part of the above diagram commutes.

It follows from the functorial properties of the theories involved that this diagram is compatible with diffeomorphisms of X; in particular all maps are W-maps.

Let us now consider an idempotent $e \in M_k(C^\infty(X)^W)$. For any $w \in W$ we have that $w^*[e] = [w^*e] = [e]$. For, if f is path-homotopic to e via a path e_t of idempotents, then w^*f is connected to $w^*e = e$ by the continuous path w^*e_t. Thus the K-class of e is invariant under the action of W. In the odd case, the class of an invertible matrix $g \in GL_k(C^\infty(X)^W)$ constist of elements of this group which are homotopic to g. But the action of W sends invertible elements to invertible elements and if h is connected to g by a continuous path of invertibles, then the same is true about $w^*g = g$ and w^*h.

By functoriality, the image of $K_*(C^\infty(X)^W)$ under the Chern character map in each of the three cases must therefore be a submodule of the invariant part of the corresponding cohomology theory on the right. In other words, we have the following commutative diagram.

$$\begin{tikzcd}
K_*(C^\infty(X)^W) \arrow[r, "ch_{CQ}"] \arrow[d, equal] & HP_*(C^\infty(X))^W \arrow[d, "\mu_c"] \\
K_*(C^\infty(X)^W) \arrow[r, "ch_{CW}"] \arrow[d, equal] & H^*_{dR}(X)^W \\
K_*(C^\infty(X)^W) \arrow[r, "ch_G"] & H^*_{AS}(X)^W \arrow[u, "\lambda"]
\end{tikzcd}$$

The two vertical arrows on the right remain isomorphisms, since the original isomorphisms were compatible with the action of the group W.

Since $C^\infty(X)^W$ holomorphically closed and dense in $C(X)^W$ the K-

theory of the two algebras is the same

$$K_*(C^\infty(X)^W) \simeq K_*(C(X)^W).$$

Furthermore, the isomorphism of C^*-algebras $C(X)^W = C(X/W)$ gives

$$K_*(C(X)^W) = K_*(C(X/W) = K^*(X/W),$$

where the last identity follows from the Serre-Swan theorem using the fact that X/W is a compact Hausdorff space. Thus we have established the isomorphism

$$K_*(C^\infty(X/W)) = K^*(X/W).$$

On the side of homology it is well known that $H^*_{AS}(X)^W = H^*_{AS}(X/W)$. On the other hand the result of Wassermann [23, p. 238] gives that $HP_*(C^\infty(X)^W) = HP_*(C^\infty(X))^W$. We thus have the following commutative diagram

$$\begin{tikzcd}
K_*(C^\infty(X)^W) \arrow[r, "ch_{CQ}"] \arrow[d, "\simeq"'] & HP_*(C^\infty(X)^W) \arrow[d, "\lambda^{-1}\circ\mu_c"'] \\
K^*(X/W) \arrow[r, "ch_G"] & H^*_{AS}(X/W, \mathbf{C})
\end{tikzcd}$$

in which the bottom map is the Chern character constructed by Gorokhovsky. Since Gorokhovsky's Chern character is the same as the classical Chern character, the bottom map becomes an isomorphism after the left hand side has been tensored with $\mathbf{C}$. Applying this reasoning to the situation described by the diagram 3.6 we see that the bottom arrow in that diagram is an isomorphism after tensoring with $\mathbf{C}$.

We have therefore proved that for every compact open subgroup K the map

$$ch^K_{CQ} : K_*(C^*_r(G//K)) \otimes_{\mathbf{Z}} \mathbf{C} \to HP_*(S(G//K))$$

is an isomorphism. Passing to the limit we finish the proof of the Theorem.

■

3.8 Remark. The result stated in Theorem 3.3 is crucial to the proof of the Baum-Connes conjecture given in [2].

References

[1] M. F. Atiyah, K-theory, Benjamin, New York, 1967.

[2] P. Baum, N. Higson, R. J. Plymen, A proof of the Baum-Connes conjecture for p-adic $GL(n)$, *C. R. Acad. Sci. Paris* **325** (1997) 171-176.

[3] J. Brodzki, R. J. Plymen, Periodic cyclic homology of certain nuclear algebras, *C. R. Acad. Sci. Paris* **329** (1999), 671-676.

[4] A. Connes, Noncommutative differential geometry, *Publ. Mathem. I.H.E.S.* **62** (1985), 257-360.

[5] A. Connes, Noncommutative geometry, Academic Press, London, New York 1994.

[6] A. Connes, H. Moscovici, Cyclic cohomology, the Novikov conjecture and hyperbolic groups, *Topology* **29**, (1990), 345-388.

[7] J. Cuntz, Bivariante K-theorie für lokcalconvexe Algebren und der Chern-Connes-Charakter, *Doc. Math* **2** (1997), 139-182.

[8] J. Cuntz, D. Quillen, Algebra extensions and nonsingularity, *Journal AMS* **8** (1995), 251-289.

[9] J. Cuntz, D. Quillen, Operators on noncommutative differential forms and cyclic homology, In: Geometry, Topology and Physics for Raoul Bott, International Press, Cambridge MA, 1995.

[10] J. Cuntz, D. Quillen, Cyclic homology and nonsingularity, *Journal AMS*, **8**, 373-442.

[11] E. Getzler, The odd Chern character in cyclic homology and spectral flow, *Topology* **33** (1994), 663-681.

[12] A. Gorokhovsky, Chern classes in Alexander-Spanier Cohomology, *K-theory* **15** (1998), 253-268.

[13] Harish-Chandra, Collected Papers, Vol. 4, Springer Verlag, Berlin 1984.

[14] J.D.S. Jones, C. Kassel, Bivariant cyclic theory, *K-theory* **3** (1989), 339-366.

[15] C. Kassel, Caractére de Chern bivariant, *K-theory* **3** (1989), 367-400.

[16] M. Karoubi, Le isomorphismes de Chern et de Thom-Ghysin en K-théorie, Séminaire Cartan-Schwartz, 16e année, 1963/64, n. 16.

[17] J.-L. Loday , Cyclic homology, Springer-Verlag, Berlin, 1992.

[18] P. Mischenko, Invariant tempered distributions on the reductive p-adic group $GL_n(F_p)$, C. R. Math. Rep. Acad. Sci. Canada 4 (1982) 123-127

[19] V. Nistor, *A bivariant Chern-Connes character*, Annals of Math. **138** (1993), 555-590.

[20] N.C. Phillips, K-theory for Fréchet algebras, *Int. J. Mathem.* **2** (1991), 77-129.

[21] D. Quillen, Superconnection character forms and the Cayley transform, *Topology* **27** (1988), 211-238.

[22] J. Rosenberg, Algebraic K-theory and its applications, GTM 147, Springer Verlag, Berlin, New York 1994.

[23] A. Wassermann, Cyclic cohomology of algebras of smooth functions on orbifolds, London Math. Soc. Lecture Notes 135 (1988) 229-244

J.B.: School of Mathematical Sciences, University of Exeter, Exeter EX4 4QE; brodzki@maths.ex.ac.uk
R.P.: Department of Mathematics, University of Manchester, Manchester M13 9PL; roger@ma.man.ac.uk